

Innovation. Quality. Solutions.



Tribunil contains 700 g/kg methabenzthiazuron (MOA Group C)

Features

- Early post-emergence weed control in onions
- Control of annual broadleaf weed seedlings
- Group C Herbicide

Background

Tribunil® Selective Herbicide has provided reliable control of a range of weeds in Australian onion crops for many years. The product was originally developed as a stand-alone herbicide, however its use has been substantially modified by grower experience. Tribunil today forms part of a herbicide program based on reduced rates that improve crop safety while still maintaining an acceptable level of weed control.

Mode of Action

Tribunil is a member of the urea group of herbicides, the mode of action of which inhibits photosynthesis at photosystem II. This means that treated weeds cannot fix CO₂ or produce the nutrients needed for their survival.

Use rates and timing

Tribunil is registered at a range of use rates, the choice of which depends on several factors including the number of true leaves present on the onion at the time of application, environmental conditions and sowing time. Soil type should also be considered and lower rates of Tribunil should be used on lighter textured soils.

Tribunil should be applied to onions once they have one or more true leaves. Spring sown crops in southern Victoria and Tasmania can be sprayed from the one leaf stage, however in other areas and when weather is warm to hot, crops should not be sprayed until the second leaf is obvious.

Weeds should be actively growing when sprayed, however if droughted, frosted or waterlogged, spraying should be delayed. When rain is expected spraying should also be delayed, as rain within 12-24 hours after spraying may reduce weed control. Crops should not be sprayed if frosted or wet with dew or rain. Always read and follow the product label.

The information provided herein may include extracts from the product label and does not constitute the complete directions for use. READ THE PRODUCT LABEL THOROUGHLY BEFORE OPENING OR USING TRIBUNIL

AgNova Technologies Pty Ltd
ABN 70 097 705 158
Unit 4, 482 Kingsford Smith Drive
Hamilton, Qld 4007 Australia
Phone (03) 9899 8100
Email info@agnova.com.au

Data referred to herein were generated using products and rates that were registered at the time. Always read and follow product labels. AgNova Technologies Pty Ltd shall not be liable for any consequential or other loss or damage relating to the supply or subsequent handling or use of this product, unless such liability by law cannot be lawfully excluded or limited. All warranties, conditions or rights implied by statute or other law which may be lawfully excluded are so excluded. Where the liability of AgNova Technologies Pty Ltd for breach of any such statutory warranties and conditions cannot be lawfully excluded but may be limited to it re-supplying the product or an equivalent product or the cost of a product or an equivalent product, then the liability of AgNova Technologies Pty Ltd for any breach of such statutory warranty or condition is so limited.
© Copyright AgNova Technologies 2018. TRBAGN210601
© Registered trademark of AgNova Technologies Pty Ltd.

Crop safety

Tribunil contains the active ingredient methabenzthiazuron and can damage onion crops under a range of conditions. Onions subjected to sand blasting, water stress or other causes can be particularly vulnerable. Herbicide damaged onion crops may be more susceptible to disease and yield less than undamaged crops.

In addition, newer onion varieties, and especially hybrids and reds, may be more sensitive to Tribunil, particularly at higher rates. Small-scale crop safety tests should be conducted with Tribunil prior to treating large areas.

Susceptible crops such as crucifers, cucurbits, lettuce or tomatoes should not be sown within 18 months after applying more than 2 kg Tribunil per hectare in one season.

Compatibility

Baron® 400 WG Selective Herbicide may be tank mixed with Tribunil according to the Directions for Use table. Other tank mixtures have not been tested and may cause severe phytotoxicity to onions. Prior to mixing Tribunil with any other product, small scale crop safety tests should be conducted prior to treating large areas. Do not add wetting agent to Tribunil.

Withholding period

Onion crops treated with Tribunil should not be harvested for seven (7) weeks after application.

Summary

- Early application from the 1 leaf stage (see product label for details)
- Wide spectrum broadleaf weed control
- Trusted & proven performance & crop safety

