

DuPont™ Linuron DF® herbicide

Technical Information

Active Constituent:
500 g/kg LINURON

Pack Sizes:
2 kg



READ SAFETY DIRECTIONS BEFORE OPENING OR USING

For the selective weed control in carrots, coriander, parsnips, cereals, onions, potatoes, soybeans, sweet corn and maize as per Directions for Use table.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with the eyes and skin. **DO NOT** inhale dust or spray mist. If product on skin, immediately wash the area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet.

GENERAL INSTRUCTIONS

This product is a foliar absorbed and residual herbicide i.e. it controls weeds present at application by absorption through the foliage and subsequent germination of weeds through root absorption. For best results apply to well cultivated, weed-free surface which is free of trash and clods. **DO NOT** cultivate after application of the product as weed control may be reduced and/or crop injury may result.

RESISTANT WEEDS WARNING

DuPont™ Linuron DF® herbicide is a member of the Urea group of herbicides. DuPont™ Linuron DF® herbicide has the inhibitors of photosynthesis at photosystem II mode of action. For weed resistance management DuPont™ Linuron DF® herbicide is a Group C herbicide.

Some naturally-occurring weed biotypes resistant to DuPont™ Linuron DF® herbicide and other inhibitors of photosynthesis at photosystem II herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds may not be controlled by DuPont™ Linuron DF® herbicide or other inhibitors of photosynthesis at photosystem II herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, DuPont accepts no responsibility for any losses that may result from the failure of DuPont™ Linuron DF® herbicide to control resistant weeds.

Strategies to minimise the risk of herbicide resistance are available. Consult your farm chemical supplier, consultants, local Department of Agriculture or Primary Industries, or local Du Pont (Australia) Ltd representative.

EQUIPMENT AND MIXING INSTRUCTIONS

DuPont™ Linuron DF® herbicide is a dry flowable formulation to be mixed with water and applied as a spray. Partially fill the spray tank with water. Measure the amount of product required as indicated in the directions for use table. Add this to the spray tank with the agitation system engaged. Top up tank to the correct water volume. To prevent nozzle blockage, strainer and nozzle screens should be 50 mesh or coarser. **THE MATERIAL MUST BE KEPT IN SUSPENSION AT ALL TIMES BY CONTINUOUS AGITATION.**

Water Volume & Spray Pressure

- Pre-emergence – 300 –500 L/ha and 200-300 kPa pressure.
- Post-emergence – Use minimum of 300 L/ha and 140-170 kPa pressure.
- Band Treatments – Use proportionately less.
- Wheat, Barley, Oats only
Aerial application – use a minimum of 20 L/ha spray under optimum conditions otherwise a minimum of 30 L/ha.
Ground application – apply in a minimum of 90 L/ha (spray).

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Product is slightly toxic to fish. **DO NOT** contaminate streams, rivers, waterways with chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, well-ventilated area, as cool as possible out of direct sunlight. **DO NOT** store for prolong periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilizers. Empty all bag contents into spray tank. **DO NOT** dispose of

undiluted chemical on site. Break, crush, puncture or bury empty containers in a local authority landfill. If not available bury the container below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and where roots of desirable plants may extend. Empty containers and product should NOT be burnt.

**IN A MEDICAL EMERGENCY CALL
1800 674 415 All hours**

NOTICE TO THE BUYER

To the extent permitted by law all conditions and warranties and statutory or other rights of action which buyer or any other user

may have against DuPont or Seller are hereby excluded. DuPont hereby give notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information or advice provided by DuPont or on its behalf unless it is established that such information or advice was provided negligently and that the product has been used strictly as directed. DuPont's liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price paid therefore.

APVMA Approval Number:60222/1005

DIRECTIONS FOR USE

RESTRAINTS:

Sufficient moisture, either by irrigation or rainfall is required within 3 - 4 days of application to carry the product into the soil profile.

Crops	Weeds Controlled	State	Rate / ha		Critical Comments
			Light Soils	Heavy Soils	
Carrots & Parsnips	Annual Broadleaved Weeds and Grasses Such as: Amaranth, Blackberry nightshade, Capeweed, Chickweed, Common sowthistle, Corn spurry, Deadnettle, Doublegee/ Threecornered Jack (Spiny Emex), Fat-hen, Hedge mustards, Heliotrope, Marsh mallow, Paterson's curse, Pigweed, Potato weed, Redshank, Scarlet Pimpernel, Staggerweed, Stinging nettle, Summer grass (Summer grass not controlled in Qld or November to February in other states), Swinecress, Variegated thistle, Wild mustard, Wild radish, Wild turnip, Winter grass	All States	2.2kg	4.5kg	Pre-emergence – Apply immediately after planting but before crop emerges. DO NOT use on light sandy soils, low in clay or organic matter.
Carrots			1.1kg	2.2kg	Post-emergence – DO NOT apply before carrots have reached the 4 true leaf stage, weeds no more than 7 cm tall, or within 5 days of any application of any emulsifiable pesticide or wetting agent as damage to crop may result. On soils high in organic matter use post-emergence rather than pre-emergence.
Coriander Seed Crops	Amsinckia, Capeweed, Charlock, Corn Gromwell (Sheepweed, White ironweed), Deadnettle, Doublegee/Threecornered Jack(s) (Spiny Emex), Fat-hen, Hedge mustard, Indian hedge mustard, Saffron thistle, Stinging nettle, Wild radish, Wild turnip	SA, Vic, NSW, Tas only	750g	1.0 kg	DO NOT use this product on fresh coriander production. Use lower rate quoted on light soils and increase up to the higher rate for heavy clay soils or soils high in organic matter. Pre-emergence – Apply immediately after sowing. Post emergence – Apply from 2 leaf onwards. Higher rates give longer residual control. Wet soil and/or rain is a benefit soon after application.
Onions	Annual Broadleaved Weeds Such as: Black bindweed, Capeweed, Charlock, Chickweed, Clover, Common sowthistle, Corny spurry, Deadnettle, Fat-hen, Field cress, Hedge mustards, Lesser swinecress, Medics, Prickly sowthistle, Shepherd's purse, Spear thistle, Stinging nettle, Wild mustard, Wild radish, Wild turnip	Qld, NSW, Vic, SA, WA only	300 g	550 g	Post-emergence only – Apply when onions are 15 cm or more in height. Retreat when necessary.
		All States	700 g to 1 kg		Post-emergence only – Apply lowest rate when onion plants are 2-3 leaf stage and high rate from 3 leaf stage onwards. Up to three applications may be used.
Potatoes	Annual Broadleaved Weeds and Grasses such as : Amaranth, Apple-of-Peru, Blackberry nightshade, Capeweed, Chickweed, Clover, Common sowthistle, Corn spurry, Deadnettle, Doublegee/ Threecornered Jacks (Spiny Emex), Fat-hen, Hedge mustard, Pigweed, Potato weed, Redshank, Shepherd's purse, Staggerweed, Stinging Nettle, Wild mustard, Wild radish, Wild turnip, Winter grass, Winter weed, Wireweed		2.2 kg	4.5 kg	Pre-emergence – Apply after planting but before emergence of crop. Incorporate by irrigation or rainfall as soon as possible after application. DO NOT cultivate after application. Use higher rate during summer months.
Soybeans	Annual Broadleaved Weeds and Grass such as: Amaranth, Apple-of-Peru, Barnyard grass, Bathurst burr, Black bindweed, Blackberry nightshade, Common sowthistle, Fat-hen, Hedge mustards, Noogoora burr, Pigweed, Potato weed, Redshank, Stinking Roger, Summer grass, Turnip weed, Wild radish	Qld, NSW, Vic, WA only	2.2 kg	5.5kg	Pre-emergence – Apply within 2-3 days after sowing. Sow beans to a depth of at least 2.5 cm. Heavy rains after application may cause crop damage. DO NOT replant treated area to other crops within 3 months of application.

Crops	Weeds Controlled	State	Rate / ha		Critical Comments
			Light Soils	Heavy Soils	
Sweetcorn, Maize	Refer to weeds under Soybeans plus Stinging nettle, Wild mustard, Wild turnip. Apply where grasses are not a problem.	All States	-	2.2 to 3.5 kg	Pre-emergence – Apply as a band or broadcast treatment immediately after planting. DO NOT use on light sandy soils low in clay or organic matter. Heavy rains after application may cause crop damage.
	Refer to weeds above. Apply where Barnyard grass or Summer grass are a problem.			4.5 to 5.5 kg	
	Refer to weeds under Soybeans plus Stinging nettle, Wild mustard and Wild turnip. Apply where grasses are not a problem.		1.1 to 2.2 kg	2.2 to 4.5 kg	Post-emergence – Apply as directed band spray to each side row when crop is 30 - 40cm high. DO NOT use misting machines. Avoid spray contact with crop foliage as injury may result. DO NOT replant treated areas to other crops within 3 months of application. If weeds are present add a wetting agent at the use rate on its label.
	Refer to Weeds above. Apply Where Barnyard grass or Summer grass are a problem.		1.1 to 3.5 kg	2.2 to 5.5kg	
Wheat, Oats, Barley	Capweed, Charlock, Corn Gromwell (Sheepweed, White ironweed), Deadnettle, Doublegee/ Threecornered Jacks (Spiny Emex), Fat-hen, Hedge mustard, Indian hedge mustard, Stinging nettle, Wild radish, Wild turnip.	All States	500 g		All Weeds – for best results apply early. DO NOT apply Linuron DF [®] if weeds are greater than 15 cm in height or diameter, or crop is beyond early tillering or crop and weeds are under drought stress. Apply when weeds are less than 5 cm in height or diameter, crop is in 2 - 4 leaf stage and soil is moist enough for good growth of crop and weed seedlings.
	As above plus Amsinckia and Saffron thistle		850 g		Apply if weeds are 5 - 15 cm in height or diameter and crop is in four leaf to early tillering stage.

NOT TO BE USED FOR ANY OTHER PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHROISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

CORIANDER AND CARROTS – DO NOT HARVEST FOR 10 WEEKS AFTER APPLICATION.

ONIONS, PARSNIPS, POTATOES, WHEAT, BARLEY, OATS, MAIZE, SWEETCORN, SOYABEANS – NOT REQUIRED WHEN USED AS DIRECTED.

The information set forth herein is furnished free of charge and is based on technical data that DuPont believes to be reliable. It is intended for use by persons having technical skill at their own discretion and risk. DuPont makes no warranties, expressed or implied, and assumes no liability in connection with any of this information. Nothing herein is to be taken as license to operate under or a recommendation to infringe on any patents.

Copyright © 2008 E.I. du Pont de Nemours and Company. All rights reserved.

The DuPont Oval Logo, DuPont™, The miracles of science®, and DF® are trademarks or registered trademarks of DuPont or its affiliates.

* Non DuPont Trademark

Du Pont (Australia) Limited
7 Eden Park Drive
Macquarie Park NSW 2113
Phone: (02) 9923 6111 or 1800 257 169
<http://www.au.ag.dupont.com>